

# Cigarette smoking, disease severity and autoantibody expression in African Americans with recent-onset rheumatoid arthritis

T R Mikuls,<sup>1</sup> L B Hughes,<sup>2</sup> A O Westfall,<sup>2</sup> V M Holers,<sup>3</sup> L Parrish,<sup>3</sup> D van der Heijde,<sup>4</sup> M van Everdingen,<sup>4</sup> G S Alarcón,<sup>2</sup> D L Conn,<sup>5</sup> B Jonas,<sup>6</sup> L F Callahan,<sup>6</sup> E A Smith,<sup>7</sup> G Gilkeson,<sup>7</sup> G Howard,<sup>2</sup> L W Moreland,<sup>2,8</sup> S L Bridges Jr<sup>2</sup>

<sup>1</sup> University of Nebraska Medical Center, Nebraska Arthritis Outcomes Research Center (NAORC), and Omaha VA Medical Center, Omaha, New England, USA; <sup>2</sup> University of Alabama at Birmingham, Birmingham, Alabama, USA; <sup>3</sup> University of Colorado Health Sciences Center, Denver, Colorado, USA; <sup>4</sup> Leiden, The Netherlands; <sup>5</sup> Emory University, Atlanta, Georgia, USA; <sup>6</sup> University of North Carolina, Chapel Hill, North Carolina, USA; <sup>7</sup> Medical University of South Carolina, Charleston, South Carolina, USA; <sup>8</sup> Division of Rheumatology and Clinical Immunology, University of Pittsburgh, Pittsburgh, USA

Correspondence to: Ted R Mikuls, Department of Medicine, University of Nebraska Medical Center and Omaha VA Medical Center, 986270 Nebraska Medical Center, Omaha, NE 68198-6270, USA; tmikuls@unmc.edu

Accepted 6 January 2008  
Published Online First  
15 January 2008

## ABSTRACT

**Objective:** To examine the association of smoking with clinical and serological features in African Americans with recent-onset rheumatoid arthritis (RA) and to explore whether this association is dependent on the presence of the *HLA-DRB1* shared epitope (SE).

**Methods:** In African Americans with recent-onset RA (n = 300), we examined the association of cigarette smoking (current versus past versus never and pack-years of exposure) with anti-cyclic citrullinated peptide antibody, rheumatoid factor (RF) (IgM and IgA), rheumatoid nodules and baseline radiographic erosions using logistic and cumulative logistic regression (adjusting for SE status). We also examined for evidence of interaction between smoking status and SE for all outcomes.

**Results:** Although there was no association with RF-IgA seropositivity, current smokers were approximately twice as likely as never smokers to have higher IgA-RF concentrations (based on tertiles; OR = 1.74; 95% CI 1.05 to 2.88) and nodules (OR = 2.43; 95% CI 1.13 to 5.22). These associations were most pronounced in those with more than 20 pack-years of exposure. There was no association of smoking status or cumulative tobacco exposure with anti-cyclic citrullinated peptide antibody, IgM-RF or radiographic erosions. There was also no evidence of a biological or statistical SE-smoking interaction for any of the outcomes examined.

**Conclusions:** This is the first study to systematically examine the association of cigarette smoking with RA-related features in African Americans. Cigarette smoking is associated with both subcutaneous nodules and higher serum concentrations of IgA-RF in African Americans with RA, associations that may have important implications for long-term outcomes in this population.

increased risk of anti-cyclic citrullinated peptide (anti-CCP) antibody-positive RA among those with a single SE-containing *HLA-DRB1* allele and a more than 20-fold increased risk in those with two SE-containing alleles.<sup>2</sup>

In addition to its association with disease risk, smoking also appears to be an important determinant of RA disease expression. Specifically, smoking has been associated with the presence of extra-articular disease manifestations<sup>14</sup> (including rheumatoid nodules), radiographic severity,<sup>15–17</sup> and serum rheumatoid factor (RF).<sup>15 16 18 19</sup> Reports examining associations of smoking with anti-CCP antibody in RA have been inconsistent with at least one study showing a positive association<sup>20</sup> while another showed no such association.<sup>21</sup> Studies to date examining the association of smoking with RA disease features have almost exclusively involved subjects of European ancestry. The lack of such studies among ethnic/racial minorities is relevant for three reasons: (a) the prevalence of smoking in African Americans appears to be increasing<sup>22</sup> while “quit rates” in African Americans are lower compared to Caucasians;<sup>23</sup> (b) disease characteristics related to smoking portend a poor outcome in RA;<sup>24–30</sup> and (c) smoking appears to have a disproportionately negative impact on minority patients with select chronic conditions.<sup>31 32</sup> The objective of this cross-sectional study was to examine the association of cigarette smoking with disease expression in a group of well characterised African Americans with recent-onset RA.

## PATIENTS AND METHODS

### Study population

RA cases (n = 300) were participants in the Consortium for the Longitudinal Evaluation of African-Americans with Early Rheumatoid Arthritis (CLEAR).<sup>33–35</sup> Subjects were enrolled through one of four sites in southeast USA (University of Alabama at Birmingham, Emory University, Medical University of South Carolina, and the University of North Carolina). The study was approved by the Institutional Review Board (IRB) at each participating centre and all study subjects provided informed written consent prior to participation. Cases satisfied the American College of Rheumatology RA classification criteria,<sup>36</sup> had less than 2 years of disease duration, and self-reported African American race/ethnicity

Since the first report by Vessey *et al* nearly 20 years ago,<sup>1</sup> cigarette smoking has been shown in numerous investigations to be associated with rheumatoid arthritis (RA) susceptibility.<sup>2–10</sup> In a national study of disease-discordant monozygotic twins, smoking was associated with a 12-fold increased risk of RA,<sup>11</sup> an association that appears to be most pronounced in those with long-term exposure (>20 years duration).<sup>7 12 15</sup> The impact of smoking on RA risk also appears to be mediated by certain genetic risk factors with the *HLA-DRB1* shared epitope (SE) being the most notable. In a recent case-control study, there was no significant association of smoking with seropositive RA in cases without the SE. Ever having smoked, however, was associated with a more than sixfold

(there was no requirement for additional first- or second-degree relatives of African American ancestry). Smoking status (current, former, never) was collected at the time of enrolment and for ever smokers, pack-years of smoking served as a measure of cumulative exposure. All subjects underwent formal examination that included an assessment for the presence of subcutaneous nodules. Radiographs of the hands and wrists were obtained at baseline (defined as <2 years duration, with a mean of 13.7 months disease duration) and scored using the van der Heijde modified Sharp score.<sup>37</sup> Validated scores were available for 192 of the 300 cases under study for whom radiographs were obtained. Subjects with available scores were classified as having either erosive disease (Sharp erosion score >0) or non-erosive disease (Sharp erosion score = 0) at baseline.

### Autoantibody measurement

Autoantibody measurements, including anti-CCP antibody and RF isotypes, were performed as previously reported.<sup>33</sup> Anti-CCP (IgG) antibodies were measured in arbitrary units (U) per ml using a commercially available second generation enzyme-linked immunosorbent assay (Diastat, Axis-Shield Diagnostics Ltd, Dundee, Scotland, UK) and were considered to be positive at a cut-off value  $\geq 5$  U/ml.<sup>33</sup> RF isotypes (IgA and IgM) were measured in international units (IU) per ml and were assessed using commercially available enzyme-linked immunosorbent assay (INOVA Diagnostics Inc., San Diego, California, USA). Positive tests were defined by levels of  $\geq 10.8$  IU/ml for IgA-RF and  $\geq 9.5$  IU/ml for IgM-RF.<sup>33</sup>

### HLA-DRB1 genotyping

High resolution *HLA-DRB1* genotyping was performed as recently described with results showing a higher frequency of SE-containing alleles in African American cases compared with controls.<sup>38</sup> In brief, genotyping was completed through DNA sequencing of exon 2 using the alleleSEQR *HLA-DRB1* reagent kit and protocol (Atria Genetics, South San Francisco, California, USA). After polymerase chain reaction amplification, sequencing was performed on an ABI 377 automated sequencer (PE Applied Biosystems, Foster City, California, USA). An additional sequence reaction was performed to analyse codon 86 sequences, thus allowing for the resolution of ambiguous results for some of the exon 2 sequences. The sequences were analysed using Assign SBT3 Software (Conexio Genomics, Applecross, Perth, Western Australia), enabling assignment of genotypes based on a recent library file of *HLA-DRB1* alleles. This method detects all SE-positive alleles. When necessary, group-specific amplifications were performed and sequence analysis conducted to differentiate among alleles within a specific group.

### Statistical analysis

Dichotomous outcomes, including autoantibody status (positive versus negative), the presence of nodules and the presence of radiographic erosions were compared by smoking status using the  $\chi^2$  test in unadjusted analyses and subsequently by using multivariable logistic regression and adjusting for the presence of SE. In an additional analysis, we examined the association of smoking status with anti-CCP antibody positivity after stratifying subjects by SE status. The odds of having higher autoantibody concentrations or higher baseline erosion scores were compared among groups defined by baseline smoking status using cumulative logistic regression models (using the cumulative logit link function to generate a cumulative odds ratio (OR)<sup>39</sup>) adjusting for SE status; for this

analysis outcome variables were categorised into tertiles. The OR and 95% confidence intervals (CI) reflect the cumulative odds of having an outcome value in the highest tertile versus the low/medium tertiles or an outcome value in either the high/medium tertiles versus the lowest tertile. In secondary analyses, we also examined the association of cumulative exposure categories (never, <10, 10–20 and >20 pack-years) with the outcomes of interest. Of 300 study participants, 14 subjects were excluded from multivariable analyses due to missing values for SE (n = 10) or smoking status (n = 4). Additional subjects with missing data for nodules (n = 2) and erosions (n = 94) were excluded from multivariable analyses examining the association of smoking with these outcomes. There were an additional six subjects with a history of smoking but missing cumulative exposure data excluded from analyses examining the association of smoking pack-years with the outcomes of interest.

Gene–environment interactions (SE–smoking) were examined in two ways, as reviewed elsewhere.<sup>40</sup> Based on recent reports of disease susceptibility,<sup>2, 6, 41</sup> we assessed the biological (additive) interaction between ever smoking (current or past smoking combined) and SE status by examining for evidence of departure from additivity of effects as the interaction criteria as described by Rothman *et al.*<sup>42</sup> Using this approach,<sup>40, 43</sup> we calculated the attributable proportion due to interaction (AP; where an AP = 0 corresponds to no interaction and an AP = 1.0 corresponds to “complete” interaction) and a corresponding 95% CI using the method of Hosmer and Lemeshow.<sup>44</sup> To assess for evidence of statistical (or multiplicative) interaction, we modelled the SE–smoking product terms, using a  $p < 0.05$  as the threshold for significance. All analyses were conducted using SAS v9.1 (SAS Inc., Cary, North Carolina, USA).

## RESULTS

Subject characteristics are summarised in table 1.

African Americans study subjects were comprised predominantly of women (83%) and had a mean age of approximately 51 years. Almost half of subjects were ever smokers, 42% were SE positive, 27% had baseline radiographic erosions and 14%

**Table 1** Subject characteristics of African Americans with recent onset rheumatoid arthritis (n = 300)

Characteristic*	Mean (SD) or number (%)
Female sex	246 (82%)
Age, years	50.6 (13.5)
Disease duration at baseline visit, months	12.8 (7.2)
Smoking status	
Never	142 (48%)
Past	66 (22%)
Current	88 (30%)
Ever smoked, <10 pack-years	63 (22%)
Ever smoked, 10–20 pack-years	34 (12%)
Ever smoked, >20 pack-years	51 (18%)
Shared epitope positive	123 (42%)
Anti-CCP antibody positive	186 (62%)
RF-IgA positive	202 (67%)
RF-IgM	211 (70%)
Baseline radiographic erosions	51 (27%)
Nodules	41 (14%)

CCP, cyclic citrullinated peptide; RF, rheumatoid factor.

\*Denominator as noted with exception of missing values for shared epitope status (n = 10), age (n = 2), smoking status (n = 4), pack-years of smoking (n = 6) among ever smokers, disease duration (n = 2), nodules (n = 5) and radiographic erosions (n = 108 radiographs unscanned at time of analysis).

had rheumatoid nodules. Only 23 subjects (8%) carried two SE containing alleles. Seropositivity was observed in 62% of subjects for anti-CCP antibody, 67% for IgA-RF and 70% for IgM-RF.

The associations of smoking status with autoantibody status (positive versus negative) and the presence of both nodules and erosions are summarised in table 2.

In univariate or SE-adjusted analyses, there was no association of smoking status with seropositivity for anti-CCP antibody, IgA-RF, or IgM-RF nor was there an association of smoking with the presence of radiographic erosions. While there was no risk associated with past smoking, current smokers were more than twice as likely as never smokers (OR = 2.43; 95% CI 1.13 to 5.22) to have rheumatoid nodules after adjusting for SE status. In analysis stratified by SE status, there were also no significant associations of smoking status with anti-CCP antibody positivity (table 2).

Our results were similar when examining the association of cumulative smoking exposure with the dichotomous outcomes of interest. There were no associations of smoking pack-years with autoantibody seropositivity or the presence of radiographic erosions (table 2). However, compared with never smokers, RA cases with high cumulative exposure (>20 pack-years) were significantly more likely to have rheumatoid nodules (SE-adjusted OR = 2.65; 95% CI 1.10 to 6.37) while cases with intermediate levels of exposure (10–20 pack-years) showed a trend towards higher risk (OR = 2.01; 95% CI 0.71 to 5.75).

There was no association of smoking status with higher serum concentrations of anti-CCP antibody or IgM-RF based on tertiles, nor was there an association with radiographic erosion scores (table 3). After adjusting for SE and compared with never smokers, current smokers were approximately two times more likely to have higher concentrations of IgA-RF (OR = 1.74; 95% CI 1.05 to 2.88). The risk of higher IgA-RF concentrations appeared to be most pronounced in those with more than 20 pack-years of cumulative exposure (SE-adjusted OR = 2.79; 95% CI 1.50 to 5.17) while those with low and intermediate levels of cumulative exposure displayed no such risk (table 3).

There was no evidence of statistical (multiplicative) interaction between smoking status (never, past, current) and SE for any of the outcomes examined (either dichotomous or ordered

categorical outcomes) with  $p > 0.10$  for all smoking-SE product terms (data not shown). Likewise, there was no evidence of significant biological (additive) interaction between ever smoking and SE in our models examining dichotomous outcomes. In each circumstance, the 95% CI for the AP due to interaction included zero. Results from the assessments of possible biological interactions are summarised in table 4.

## DISCUSSION

To our knowledge, this is the first study to examine the associations of cigarette smoking with RA-related disease characteristics in African Americans. In this population, current smoking is positively associated with the presence of both rheumatoid nodules and higher serum concentrations of IgA-RF, associations that appear to be independent of underlying SE status and most pronounced in those with higher levels of cumulative tobacco exposure. In contrast, our results suggest that smoking is not a major determinant of IgM-RF or anti-CCP antibody in African Americans with RA, either in terms of seropositivity or higher serum concentrations. In contrast to subjects of European ancestry with both early<sup>16</sup> and established disease,<sup>15 17</sup> smoking does not appear to be strongly associated with the presence or extent of baseline radiographic erosions among African Americans with recent-onset RA.

The association of cigarette smoking with rheumatoid nodules and higher IgA-RF concentrations may have important implications in this population. IgA-RF, for instance, has been shown in patients with RA of European/Caucasian descent to be an independent predictor of severe joint damage in established disease and has been associated with both poor functional status and extra-articular disease including pulmonary involvement.<sup>45–48</sup> In one study, 80% of patients with RA with elevated IgA-RF concentrations had one or more extra-articular disease manifestations compared with only 21% of those with increased concentrations of IgM-RF but normal IgA-RF,<sup>48</sup> highly relevant given the association of extra-articular findings with disease-related mortality.<sup>49</sup> Rheumatoid nodules, the most common extra-articular disease manifestation, have been associated with a more than fourfold increased risk of mortality in men with RA from the UK Norfolk Arthritis Registry.<sup>50</sup>

**Table 2** Associations of smoking status (current versus past versus never) and categories of cumulative smoking exposure (pack-years) with autoantibody status (positive versus negative), nodules (present versus absent) and erosive disease (present versus absent) among African Americans with recent-onset rheumatoid arthritis; all logistic regression models adjusted for presence of SE\*

	Anti-CCP positivity		RF-IgM positivity		RF-IgA positivity		Radiographic erosions		Subcutaneous nodules	
	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)	%	OR (95% CI)
Smoking status		(n = 286)		(n = 286)		(n = 286)		(n = 192)		(n = 284)
Never	64	Referent	70	Referent	67	Referent	25	Referent	10	Referent
Past	63	0.99 (0.53 to 1.85)	71	1.05 (0.54 to 2.02)	69	1.11 (0.59 to 2.10)	28	1.17 (0.52 to 2.60)	9	0.90 (0.33 to 2.47)
Current	60	0.88 (0.50 to 1.55)	70	1.05 (0.57 to 1.91)	67	1.00 (0.56 to 1.79)	27	1.11 (0.52 to 2.34)	21	2.43 (1.13 to 5.22)
Pack-years†		(n = 280)		(n = 280)		(n = 280)		(n = 189)		(n = 278)
Never	64	Referent	70	Referent	67	Referent	25	Referent	10	Referent
<10	62	0.89 (0.47 to 1.69)	70	0.97 (0.49 to 1.90)	67	0.96 (0.50 to 1.84)	23	0.89 (0.38 to 2.10)	12	1.13 (0.43 to 2.97)
10–20	58	0.83 (0.38 to 1.82)	64	0.79 (0.35 to 1.77)	61	0.78 (0.35 to 1.71)	24	0.94 (0.33 to 2.66)	18	2.01 (0.71 to 5.75)
>20	64	1.11 (0.56 to 1.36)	78	1.65 (0.76 to 3.58)	74	1.46 (0.70 to 3.02)	32	1.44 (0.60 to 3.43)	22	2.65 (1.10 to 6.37)

CCP, cyclic citrullinated peptide; RF, rheumatoid factor; SE, shared epitope.

\*Univariate p-values ( $\chi^2$  test) for smoking status:  $p = 0.75$  for anti-CCP antibody;  $p = 0.95$  for RF-IgM;  $p = 0.88$  for RF-IgA;  $p = 0.93$  for erosions;  $p = 0.02$  for nodules; Univariate p-values for categories of pack-years (Mantel-Haenszel  $\chi^2$  test):  $p = 0.73$  for anti-CCP antibody;  $p = 0.71$  for RF-IgM;  $p = 0.74$  for RF-IgA;  $p = 0.52$  for erosions;  $p = 0.03$  for nodules; in analyses stratified by SE status (0 vs 1 or 2 copies), there was no significant association of current (OR = 0.7; 95% CI 0.3 to 1.4) or former smoking (OR = 0.9; 95% CI 0.4 to 1.9) (referent to never smoking) with anti-CCP antibody status in SE-negative subjects; likewise, there was no significant association of current (OR = 1.3; 95% CI 0.5 to 3.4) or former smoking (OR = 1.3; 95% CI 0.5 to 3.5) with anti-CCP antibody status in SE-positive subjects.

†Excludes 10 subjects with missing data for SE and four subjects missing smoking data; an additional six subjects a known smoking history (three with available erosion scores) were excluded from analysis due to missing values for pack-years of smoking.

**Table 3** Associations of smoking status (current versus past versus never) and categories of cumulative smoking exposure (pack-years) with higher tertiles of autoantibody concentrations and modified Sharp erosion scores among African Americans with recent-onset rheumatoid arthritis; cumulative logistic regression models (using cumulative logit link function to generate a cumulative OR and 95% CI) adjusted for presence of SE\*

	Anti-CCP antibody OR (95% CI)	RF-IgM OR (95% CI)	RF-IgA OR (95% CI)	Erosion score OR (95% CI)
Smoking status	(n = 286)	(n = 286)	(n = 286)	(n = 192)
Never	Referent	Referent	Referent	Referent
Past	1.13 (0.65 to 1.94)	1.15 (0.67 to 1.98)	1.51 (0.88 to 2.61)	1.24 (0.56 to 2.71)
Current	1.11 (0.67 to 1.84)	1.27 (0.77 to 2.09)	1.74 (1.05 to 2.88)	1.05 (0.50 to 2.21)
Pack-years†	(n = 280)	(n = 280)	(n = 280)	(n = 189)
Never	Referent	Referent	Referent	Referent
<10	1.21 (0.69 to 2.12)	1.05 (0.60 to 1.83)	1.25 (0.71 to 2.18)	0.92 (0.40 to 2.14)
10–20	0.87 (0.43 to 1.77)	1.05 (0.52 to 2.11)	1.30 (0.64 to 2.62)	0.98 (0.35 to 2.71)
>20	1.27 (0.70 to 2.78)	1.65 (0.90 to 3.02)	2.79 (1.50 to 5.17)	1.18 (0.61 to 2.26)

CCP, cyclic citrullinated peptide; RF, rheumatoid factor; SE, shared epitope.

\*Outcome variables categorised into tertiles; OR (and 95% CI) reflects the cumulative odds of having an outcome value in the highest tertile versus the low/medium tertiles or an outcome value in either the high/medium tertiles versus the lowest tertile.

†Six subjects with a known smoking history (three with available erosion scores) were excluded from analysis due to missing values for pack-years of smoking.

It remains unknown whether smoking is a risk factor in African Americans for the development of RA (disease characterised by the presence of higher IgA-RF values) or whether smoking or its by-products stimulate specific autoantibody expression. In a cross-sectional study of subjects of European ancestry without RA, both IgA-RF (34.4%) and IgM-RF (34.1%) were significantly more likely to be positive in smokers compared with non-smokers (21.9%), suggesting that smoking may have a direct effect on humoral immunity irrespective of disease status.<sup>51</sup> Padyukov *et al* found that smoking increases the risk of developing RF-positive RA but not seronegative RA, a risk that was confined to those positive for SE.<sup>6</sup> Although Padyukov and colleagues did not examine specific RF isotypes or anti-CCP antibody, their results support the existence of SE–smoking interaction in the risk of RF-positive disease. In a more recent study, Klareskog *et al* found a similar association of smoking for the development of anti-CCP antibody positive disease, a risk also limited to SE-positive subjects.<sup>2</sup> Based on subanalyses in subjects discordant for anti-CCP antibody and RF, their results also suggest that the development of anticitrulline immunity is the “primary” pathogenic event in smokers carrying SE while the development of RF more likely represents a “secondary” phenomenon.

In contrast to recent reports examining disease susceptibility<sup>2 6 41</sup> in Caucasians and the effect of smoking on autoantibody expression,<sup>20</sup> we found no evidence to support the role of a

SE–smoking interaction in RA disease expression among African Americans. However, our results are consistent with recent findings from three large North American Caucasian RA cohorts, showing no major SE–smoking interaction for anti-CCP antibody expression.<sup>52</sup> In a report of 407 Dutch subjects with recent-onset disease, a history of tobacco exposure was associated with an increased odds of anti-CCP antibody positivity in SE-positive RA subjects but not in those negative for SE.<sup>20</sup> In contrast to our study, biological interaction between SE and smoking in the study of Linn-Rasker *et al*<sup>20</sup> was defined as being present because the odds of having anti-CCP antibody for subjects having both tobacco exposure and SE (OR = 5.27; 95% CI 2.37 to 11.80) was higher than the summed odds ratios of subjects with only tobacco exposure (OR = 1.07; 95% CI 0.43 to 2.65) or SE alone (OR = 2.49; 95% CI 1.18 to 5.31). In post-hoc analyses of these data, there was no evidence of statistical interaction (p-value of interaction term = 0.2) while the AP due to interaction was estimated to be 0.5<sup>53 54</sup>. Even with the use of the less stringent definition,<sup>20</sup> our results still provide no conclusive evidence of an interaction between smoking and SE for the development of anti-CCP antibody, RF (either IgM or IgA isotype), or radiographic erosions in African Americans with RA (although suggesting the presence of a borderline SE–smoking interaction for the presence of nodules).

Although our study includes the largest African American RA cohort systematically examined to date, it is likely that our

**Table 4** Assessment of biological (additive) interaction between ever smoking and HLA-DRB1 SE status among African Americans with recent-onset rheumatoid arthritis; assessed using departure from additivity of effects as the interaction criteria\*

	Never smoker/SE-negative	Never smoker/SE-positive OR (95% CI)	Ever smoker/SE-negative OR (95% CI)	Ever smoker/SE-positive OR (95% CI)	Attributable proportion due to interaction (95% CI)
Anti-CCP positivity (n = 286)	Referent	1.74 (0.85 to 3.55)	0.76 (0.41 to 1.41)	2.23 (1.07 to 4.67)	0.33 (−0.30 to 0.96)
RF-IgM positivity (n = 286)	Referent	1.85 (0.87 to 3.95)	0.96 (0.51 to 1.83)	2.25 (1.02 to 4.94)	0.44 (−0.57 to 0.95)
RF-IgA positivity (n = 286)	Referent	1.51 (0.73 to 3.13)	1.02 (0.54 to 1.93)	1.64 (0.79 to 3.43)	0.07 (−0.77 to 0.91)
Erosive status (n = 192)	Referent	0.91 (0.34 to 2.41)	0.97 (0.40 to 2.33)	1.24 (0.50 to 3.08)	0.29 (−0.69 to 1.00)
Nodules (n = 284)	Referent	0.94 (0.31 to 2.89)	1.23 (0.47 to 3.24)	2.30 (0.89 to 5.98)	0.49 (−0.17 to 1.00)

CCP, cyclic citrullinated peptide; RF, rheumatoid factor; SE, shared epitope.

\*No evidence of interaction when attributable proportion = 0.

analysis was not adequately powered to detect this gene-environment interaction. Because these subjects were recruited primarily from academic rheumatology practices, it is possible that these results are not generalisable to other African American populations with RA. Given the lower prevalence of *HLA-DRB1* alleles in African American cases relative to Caucasians with RA,<sup>55</sup> we were not able to adequately examine the impact of SE “dose” nor were we able to fully evaluate the risk conferred by specific *HLA-DRB1* alleles. This limitation is noteworthy because different SE alleles appear to differ in their interaction with smoking and predisposition to autoantibodies in patients of European ancestry.<sup>56</sup> It is possible that discrepant results observed across different populations (African American versus Caucasian and/or North American versus European) may be due to other “unmeasured” risk factors involved in RA pathogenesis that are more common to select ethnic/racial groups (i.e. additional genes and/or environmental triggers).

In summary, the data presented in this report show that cigarette smoking is associated with select disease characteristics in African Americans with recent onset RA, features that include higher serum concentrations of IgA-RF and the presence of subcutaneous nodules. As reported in other studies of disease susceptibility, the associated risk of smoking appears to be greatest among subjects with the highest levels of cumulative exposure. Further studies that include a larger number of well-characterised cases and controls with extended follow-up will be needed to better define the complex relationship of smoking with predisposing risk alleles in RA susceptibility and disease expression in African Americans.

**Acknowledgements:** The CLEAR Registry is supported by NIH grant N01-AR-02247. TRM's work was supported by grants from NIH/NIAMS (R03-AR-054539) and K23-AR-050004) and the Arthritis Foundation (National and Nebraska Chapters). The CLEAR Registry is also supported by NIH N01-AR-6-2278 (SLB, PI) and by the UAB GCRC and grant from NIH/NCRC (M01-RR-00032).

**Competing interests:** None.

## REFERENCES

- Vessey MP, Villard-Mackintosh L, Yeates D. Oral contraceptives, cigarette smoking and other factors in relation to arthritis. *Contraception* 1987;**35**:457–64.
- Klareskog L, Stolt P, Lundberg K, Kallberg H, Bengtsson C, Grunewald J, et al. A new model for an etiology of rheumatoid arthritis: smoking may trigger HLA-DR (shared epitope)-restricted immune reactions to autoantigens modified by citrullination. *Arthritis Rheum* 2006;**54**:38–46.
- Karlson EW, Lee IM, Cook NR, Manson JE, Buring JE, Hennekens CH. A retrospective cohort study of cigarette smoking and risk of rheumatoid arthritis in female health professionals. *Arthritis Rheum* 1999;**42**:910–17.
- Heliövaara M, Aho K, Aromaa A, Knekt P, Reunanen A. Smoking and risk of rheumatoid arthritis. *J Rheumatol* 1993;**20**:1830–5.
- Symmons DP, Bankhead CR, Harrison BJ, Brennan P, Barrett EM, Scott DG, et al. Blood transfusion, smoking, and obesity as risk factors for the development of rheumatoid arthritis: results from a primary care-based incident case-control study in Norfolk, England. *Arthritis Rheum* 1997;**40**:1955–61.
- Padyukov L, Silva C, Stolt P, Alfredsson L, Klareskog L. A gene-environment interaction between smoking and shared epitope genes in HLA-DR provides a high risk of seropositive rheumatoid arthritis. *Arthritis Rheum* 2004;**50**:3085–92.
- Criswell LA, Merlino LA, Cerhan JR, Mikuls TR, Mudano AS, Burma M, et al. Cigarette smoking and the risk of rheumatoid arthritis among postmenopausal women: results from the Iowa Women's Health Study. *Am J Med* 2002;**15**:465–71.
- Silman AJ, Newman J, MacGregor AJ. Cigarette smoking increases the risk of rheumatoid arthritis. *Arthritis Rheum* 1996;**39**:732–5.
- Voigt LF, Koepsell TD, Nelson JL, Dugowson CE, Daling JR. Smoking, obesity, alcohol consumption, and the risk of rheumatoid arthritis. *Epidemiology* 1994;**5**:525–32.
- Uhlig T, Hagen KB, Kvien TK. Current tobacco smoking, formal education, and the risk of rheumatoid arthritis. *J Rheumatol* 1999;**26**:47–54.
- Silman A, Newman J, MacGregor A. Cigarette smoking increases the risk of rheumatoid arthritis. *Arthritis Rheum* 1996;**39**:732–5.
- Costenbader K, Feskanich D, Mandl L, Karlson EW. Smoking intensity, duration, and cessation, and the risk of rheumatoid arthritis in women. *Am J Med* 2006;**119**:503.e501–9.
- Stolt P, Bengtsson C, Nordmark B, Lindblad S, Lundberg I, Klareskog L, et al. Quantification of the influence of cigarette smoking on rheumatoid arthritis: results from a population based case-control study, using incident cases. *Ann Rheum Dis* 2003;**62**:835–41.
- Turesson C, O'Fallon WM, Crowson CS, Gabriel SE, Matteson EL. Extra-articular disease manifestations in rheumatoid arthritis: induce trends and risk factors over 46 years. *Ann Rheum Dis* 2003;**62**:722–7.
- Mattey DL, Hutchinson D, Dawes PT, Nixon NB, Clarke S, Fisher J, et al. Smoking and disease severity in rheumatoid arthritis: association with polymorphism at the glutathione S-transferase M1 locus. *Arthritis Rheum* 2002;**46**:640–6.
- Papadopoulos NG, Alamanos Y, Voulgari PV, Epagelis EK, Tsigetaki N, Drosos AA. Does cigarette smoking influence disease expression, activity, and severity in early rheumatoid arthritis patients? *Clin Exp Rheumatol* 2005;**23**:861–6.
- Saag KG, Cerhan JR, Kolluri S, Ohashi KO, Hunninghake GW, Schwartz DA. The effects of cigarette smoking on rheumatoid arthritis disease severity. *Ann Rheum Dis* 1997;**56**:463–9.
- Wolfe F. The effect of smoking on clinical, laboratory, and radiographic status in rheumatoid arthritis. *J Rheumatol* 2000;**27**:630–7.
- Harrison BJ, Silman AJ, Wiles NJ, Scott DG, Symmons DP. The association of cigarette smoking with disease outcome in patients with early inflammatory polyarthritis. *Arthritis Rheum* 2001;**44**:323–30.
- Linn-Rasker SP, van der Helm-van Mil AH, van Gaalen FA, Kloppenburg M, de Vries R, le Cessie S, et al. Smoking is a risk factor for anti-CCP antibodies only in RA patients that carry HLA-DRB1 shared epitope alleles. *Ann Rheum Dis* 2005;**65**:366–71.
- Jawaheer D, Lum RF, Gregersen PK, Criswell LA. Influence of male sex on disease phenotype in familial rheumatoid arthritis. *Arthritis Rheum* 2006;**54**:3087–94.
- Centers for Disease Control and Prevention. Tobacco use among high school students – United States, 1997. *MMWR* 1998;**46**:433–40.
- US Department of Health and Human Services. Tobacco use among US racial/ethnic groups – African Americans, American Indian and Alaska Natives, Asian Americans and Pacific Islanders, and Hispanics. Atlanta: US Department of Health and Human Services and Center for Disease Control and Prevention, 1998.
- Bukhari M, Lunt M, Harrison BJ, Scott DG, Symmons DP, Silman AJ. Rheumatoid factor is the major predictor of increasing severity of radiographic erosions in rheumatoid arthritis: results from the Norfolk Arthritis Register Study, a large inception cohort. *Arthritis Rheum* 2002;**46**:906–12.
- Berglin E, Johansson T, Sundin U, Jidell E, Wadell G, Hallmans G, et al. Radiological outcome in rheumatoid arthritis is predicted by the presence of antibodies against cyclic citrullinated peptide before and at disease onset, and by IgA-rheumatoid factor at disease onset. *Ann Rheum Dis* 2005;**65**:453–8.
- Heliövaara M, Aho K, Knekt P, Aromaa A, Maatela J, Reunanen A. Rheumatoid factor, chronic arthritis and mortality. *Ann Rheum Dis* 1995;**54**:811–14.
- Forslind K, Ahlmen M, Eberhardt K, Hafstrom I, Svensson B. Prediction of radiological outcome in early rheumatoid arthritis in clinical practice: role of antibodies to citrullinated peptides (anti-CCP). *Ann Rheum Dis* 2004;**63**:1090–5.
- Kastbom A, Strandberg G, Lindroos A, Skogh T. Anti-CCP antibody test predicts the diseases course during 3 years in early rheumatoid arthritis (the Swedish TIRA project). *Ann Rheum Dis* 2004;**63**:1085–9.
- Scott DL. Radiological progression in established rheumatoid arthritis. *J Rheumatol Suppl* 2004;**69**:55–65.
- Sihvonen S, Korpela M, Mustila A, Mustonen J. The predictive value of rheumatoid factor isotypes, anti-cyclic citrullinated peptide antibodies, and antineutrophil cytoplasmic antibodies for mortality in patients with rheumatoid arthritis. *J Rheumatol* 2005;**32**:2089–94.
- Haiman CA, Stram DO, Wilkens LR, Pike MC, Kolonel LN, Henderson BE, et al. Ethnic and racial differences in the smoking-related risk of lung cancer. *N Engl J Med* 2006;**354**:333–42.
- Stat bite: mortality from lung and bronchus cancer by race/ethnicity, 1998–2002. *J Natl Cancer Inst* 2006;**98**:158.
- Mikuls TR, Holers VM, Parrish LA, Conn DL, Gilkeson G, et al. Anti-cyclic citrullinated peptide antibody and rheumatoid factor isotypes in African Americans with early rheumatoid arthritis. *Arthritis Rheum* 2006;**54**:3057–9.
- Mikuls TR, Saag KG, Curtis J, Bridges SL, Alarcon GS, Westfall AO, et al. Prevalence of osteoporosis and osteopenia among African Americans with early rheumatoid arthritis: the impact of ethnic-specific normative data. *J Natl Med Assoc* 2005;**97**:1155–60.
- Bridges SL Jr, Hughes LB, Mikuls TR, Howard G, Tiwari HK, Alarcon GS, et al. Early rheumatoid arthritis in African-Americans: the CLEAR registry. *Clin Exp Rheumatol* 2003;**21**(5 Suppl 31):S138–45.
- Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper GS, et al. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arthritis Rheum* 1988;**31**:315–24.
- van der Heijde D. How to read radiographs according to the Sharp/van der Heijde method. *J Rheumatol* 1999;**26**:743–5.
- Hughes LB, Morrison D, Padilla M, Vaughn K, Westfall AO, Mikuls TR, et al. Susceptibility to rheumatoid arthritis in African Americans is associated with HLA DRB1 alleles containing shared epitope through genetic admixture with the European population. *Arthritis Rheum* 2008;**58**:349–58.
- McCullagh P, Nelder JA. *Generalized linear models*. London: Chapman & Hall, 1989.
- Ahlbom A, Alfredsson L. Interaction: a word with two meanings creates confusion. *Eur J Epidemiol* 2005;**20**:563–4.
- Kallberg H, Padyukov L, Plenge RM, Ronnelid J, Gregersen PK, van der Helm-van Mil AHM, et al. Gene-gene and gene-environment interactions involving HLA-DRB1, PTPN22, and smoking in two subsets of rheumatoid arthritis. *Am J Hum Genet* 2007;**80**:867–75.

42. **Rothman KJ**, Greenland S, Walker AM. Concepts of interaction. *Am J Epidemiol* 1980;**112**:467–70.
43. **Andersson T**, Alfredsson L, Kallberg H, Zdravkovic S, Ahlbom A. Calculating measures of biologic interaction. *Eur J Epidemiol* 2005;**20**:575–9.
44. **Hosmer DW**, Lemeshow S. Confidence interval estimation of interaction. *Epidemiology* 1992;**3**:452–6.
45. **Ates A**, Kinikli G, Turgay M, Akay G, Tokgoz G. Effects of rheumatoid factor isotypes on disease activity and severity in patients with rheumatoid arthritis: a comparative study. *Clin Rheumatol* 2007;**26**:538–45.
46. **Arinbjarnarson S**, Jonsson T, Steinsson K, Sigfusson A, Jonsson H, Geirsson A, *et al.* IgA rheumatoid factor correlates with changes in B and T lymphocyte subsets and disease manifestations in rheumatoid arthritis. *J Rheumatol* 1997;**24**:269–74.
47. **Jonsson T**, Thorsteinsson H, Arinbjarnarson S, Thorsteinsson J, Valdimarsson H. Clinical implications of IgA rheumatoid factor subclasses. *Ann Rheum Dis* 1995;**54**:578–81.
48. **Jonsson T**, Arinbjarnarson S, Thorsteinsson J, Steinsson K, Geirsson AJ, Jonsson H, *et al.* Raised IgA rheumatoid factor (RF) but not IgM RF is associated with extra-articular manifestations in rheumatoid arthritis. *Scand J Rheumatol* 1995;**24**:372–5.
49. **Gabriel SE**, Crowson CS, Kremers HM, Doran MF, Turesson C, O'Fallon WM, *et al.* Survival in rheumatoid arthritis: a population-based analysis of trends over 40 years. *Arthritis Rheum* 2003;**48**:54–8.
50. **Goodson NJ**, Wiles NJ, Lunt M, Barrett EM, Silman AJ, Symmons DPM. Mortality in early inflammatory polyarthritis: cardiovascular mortality is increased in seropositive patients. *Arthritis Rheum* 2002;**46**:2010–19.
51. **Jonsson T**, Thorsteinsson J, Valdimarsson H. Does smoking stimulate rheumatoid factor production in non-rheumatic individuals? *APMIS* 1998;**106**:970–4.
52. **Lee H**, Irigoyen P, Kern M, Lee A, Batliwalla F, Khalili H, *et al.* Interaction between smoking, the shared epitope, and anti-cyclic citrullinated peptide: a mixed picture in three large North American rheumatoid arthritis cohorts. *Arthritis Rheum* 2007;**56**:1745–53.
53. **Mattey DL**. No association of smoking with anti-CCP antibodies in RA after adjustment for rheumatoid factor. *Ann Rheum Dis* 2005;**eLetter**:(25 August 2005).
54. **Costenbader K**, Chibnik L, Mandl L, Karlson EW. Testing for gene-environment interaction. *Ann Rheum Dis* 2006;**eLetter**:(6 January 2006).
55. **Gorman JD**, Lum RF, Chen JJ, Suarez-Almazor ME, Thomson G, Criswell LA. Impact of shared epitope genotype and ethnicity on erosive disease: a meta-analysis of 3,240 rheumatoid arthritis patients. *Arthritis Rheum* 2004;**50**:400–12.
56. **van der Helm-van Mil AH**, Verpoort KN, le Cessie S, Huizinga TW, de Vries RR, Toes RE. The HLA-DRB1 shared epitope alleles differ in the interaction with smoking and predisposition to antibodies to cyclic citrullinated peptide. *Arthritis Rheum* 2007;**56**:425–32.

### Need a helping hand with your career choices?

If you need to take stock, get some career advice or find out about the choices available to you then the **BMJ Careers Fair** is the place to do it. You can find out about how best to present yourself to potential employers, polishing up your CV, working abroad, locum working and much more.

#### BMJ Careers Fairs – dates for your diary

3–4 October 2008 – Business Design Centre, London

10–11 October 2008 – Thinktank, The Science Museum, Birmingham – working in partnership with the West Midlands Deanery

Register now at [bmjcareersfair.com](http://bmjcareersfair.com)

**BMJ Careers**